

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

Effect of dietary inclusion of different lipid supplements on quality and oxidative susceptibility of beef meat

This is the author's manuscript

Original Citation:

Availability:

This version is available <http://hdl.handle.net/2318/1689142> since 2020-01-14T12:01:24Z

Published version:

DOI:10.1080/1828051X.2018.1485517

Terms of use:

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



Effect of dietary inclusion of different lipid supplements on quality and oxidative susceptibility of beef meat

Journal:	<i>Italian Journal of Animal Science</i>
Manuscript ID	TJAS-2016-0320.R4
Manuscript Type:	Short Communication
Date Submitted by the Author:	n/a
Complete List of Authors:	Mordenti, Attilio; Università degli Studi di Bologna, DIMEVET Brogna, Nico; Università degli Studi di Bologna, DIMEVET Merendi, Flavia; Università degli Studi di Bologna, DIMEVET Formigoni, Andrea; Università degli Studi di Bologna, DIMEVET Sardi, Luca; Università degli Studi di Bologna Dipartimento di Scienze Mediche Veterinarie Cardenia, Vladimiro; Università degli Studi di Bologna, DISTAL Rodriguez Estrada, Maria; Università degli Studi di Bologna, DISTAL
Keywords:	beef quality, flaxseed, CLA, vitamin E, lipid fatty acid composition and oxidation
Abstract:	The aim of this study was to evaluate the impact of dietary extruded flaxseed (EF) and/or rumen-protected (rp) lipids on beef meat quality and oxidation. Sixty-three crossbred heifers (Charolais X Limousin) were evenly distributed into seven experimental groups, balanced in terms of age and BW. The feeding groups differed in both, the dietary lipid source (EF and/or rp conjugated linoleic acid (CLA), supplemented with vitamin E (VE)) and the supplementation length (90 or 180 days before slaughter); the same total amount of lipids was administered to the animals. With the respect of the control group, the α -linolenic acid content significantly increased and the n-6/n-3 ratio decreased in cattle groups fed with EF. The results also

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

	show that providing CLA and/or EF did not lead to an increase in CLA isomers (CLAs) in the lipid fraction of the meat. Our finding show that administering EF for a shorter period with respect to longer, moderately enhance the nutritional value of the meat with regard to fat composition. No significant effects of dietary supplementation on TBA-reactive substances (TBARS) levels of fresh meat were observed; however, the protection provided by vitamin supplementation might become evident during meat storage under commercial retail conditions.

SCHOLARONE™
Manuscripts

For Peer Review Only

Effect of dietary inclusion of different lipid supplements on quality and oxidative susceptibility of beef meat

A.L. Mordenti^{1*}, N. Brogna¹, F. Merendi¹, A. Formigoni¹, L. Sardi¹, V. Cardenia², M.T. Rodriguez-Estrada²

¹ *Dipartimento di Scienze Mediche Veterinarie, Università di Bologna (DIMEVET; Via Tolara di sopra 50; 40064, Ozzano Emilia, Bologna), Italy.*

² *Dipartimento di Scienze e Tecnologie Agro-Alimentari, Università di Bologna (DISTAL; Viale G. Fanin 40; 40127, Bologna), Italy.*

*Corresponding author: Attilio Luigi Mordenti; <http://orcid.org/0000-0002-4162-6726>; Phone: +390512097381; Fax: +390512097373; email: attilio.mordenti@unibo.it

Keywords: beef quality; flaxseed; CLA; vitamin E; lipid fatty acid composition and oxidation.

Abstract

The aim of this study was to evaluate the impact of dietary extruded flaxseed (EF) and/or rumen-protected (rp) lipids on beef meat quality and oxidation. Sixty-three crossbred heifers (Charolais X Limousin) were evenly distributed into seven experimental groups, balanced in terms of age and BW. The feeding groups differed in both, the dietary lipid source (EF and/or rp conjugated linoleic acid (CLA), supplemented with vitamin E (VE)) and the supplementation length (90 or 180 days before slaughter); the same total amount of lipids was administered to the animals. With the respect of the control group, the α -linolenic acid content significantly increased and the n-6/n-3 ratio decreased in cattle groups fed with EF. The results also show that providing CLA and/or EF did not lead to an increase in CLA isomers (CLAs) in the lipid fraction of the meat. Our finding show that

administering EF for a shorter period with respect to longer, moderately enhance the nutritional value of the meat with regard to fat composition. No significant effects of dietary supplementation on TBA-reactive substances (TBARs) levels of fresh meat were observed; however, the protection provided by vitamin supplementation might become evident during meat storage under commercial retail conditions.

Introduction

In the common consumer perception of beef meat is considered a high-fat food and its consumption may increase the risk of cardiovascular disease and colon cancer in humans (McAfee et al., 2010; McNeill, 2014). Recent nutritional guidelines (EFSA, 2005; U.S. Department of Health and Human Services, 2010) recommend that people enhance their intake with foods rich in polyunsaturated fatty acids (PUFA). Particularly the intake of PUFA n-3 should be increased, so that the western diet n-6/n-3 FA ratio can decrease from 16/1 to 5/1, thus promoting health conditions and preventing the risk of diseases (Moloney et al, 2008). Human health also seems to significantly benefit from the intake of several CLAs, two of which are commonly found in beef and dairy products and are known to possess biological activity: the anti-carcinogenic and anti-atherogenic effects of cis-9, trans-11 CLA and the anti-obesity effects of trans-10, cis-12 CLA have been documented in animal experiments (McAfee et al., 2010; Salter, 2013). With the aim to positively influence the FA composition of beef intramuscular fat, pasture-based farming is the better strategy (Priolo et al., 2001). The possibility to influence the FA content can be achieved also in intensive farming conditions, with cereals and corn silage based diets, by modifying the composition of the lipid fraction of these diets (Pouzo et al., 2016; Scollan et al., 2014; Juarez et al., 2012). However, a higher content of PUFA makes meat more prone to oxidation, which can impact its overall quality and sensory traits (such as colour, flavour, aroma) (Priolo et al., 2001). This may also compromise animal health and could result in the development of unhealthy oxidation products in beef, together with decreased colour stability with a consequent reduction in shelf life (Moloney et al. 2008).

Therefore, the addition of suitable dietary protections, such as vitamin E (VE), may be important to control oxidation, especially during storage (Juarez et al., 2012). In fact, VE supplementation can lengthen the retail shelf-life of beef by 1.6 to 5 days (Nassu et al., 2011b). The aim of this study was to improve the meat nutritional properties by evaluating the impact of these dietary lipid treatments on beef meat quality (total fat content, FA composition and oxidation susceptibility).

Materials and methods

The Scientific Ethics Committee on Animal Experimentation of the University of Bologna approved the experimental protocol of this study (n.: 71674-X/6 - All.: 63). Sixty-three crossbred heifers (Charolais X Limousin), approximately 11 months old and 380 kg weight, were randomly distributed into seven groups. Each group received a basal diet included (kg/head/d): corn silage (8), beet pulp silage (5), corn meal (2.5), straw (0.8) and a concentrate (2). Feeding was enriched with various experimental supplements (EF, D α -tocopheril acetate (VE), CLAs (LUTA-CLA 60[®] (BASF) guarantees a minimum of 56% of conjugated linoleic acid methyl ester (28% cis-9, trans-11 CLA and 28% trans-10, cis-12 CLA)) at different doses, leading to 7 independent diet groups (Table 1). The diets of the experimental groups differed with respect to the lipid source and to the length of administration but, considering their entire experimental period (180 days), the same total amount of lipids was administered to the animals. When animals reached an average BW of 675 kg, they were transported to a slaughtering facility (Unipeg, Reggio Emilia). Twenty-four hours after slaughtering, a three-rib sample (20 cm of thickness from the 7th-9th thoracic vertebrae) of *Longissimus thoracis et lumborum* muscle (LT), was taken from the half right carcass of each of the 63 heifers. The muscle samples were vacuum-packaged and wet-aged for 14 days in a refrigerator at 2-4 °C. At the end of the aging period, each sample was cut into three subsamples, then vacuum-packaged, rapidly frozen and stored at -20 °C. At the time of analysis, the 63 beef samples were thawed in 3 different work sessions. Each first subsample was deboned and the first and last slices were cut and discarded. After 30 min of exposure to air, colour (L*, a*, b*) was measured using a

Minolta Chromameter CR-200 (Illuminant D65, aperture size 8 mm; Minolta Camera, Osaka, Japan). The LT muscle was then divided into a 50 x 50 x 10 mm thick sample to measure cooking loss; thereafter, in the same cooked sample, Warner-Bratzler Shear Force (WBSF) was evaluated. The WBSF method was assessed also on raw meat. Cooking loss (%) was determined using the Honikel gravimetric method (Honikel, 1998). The WBSF test was carried out on the cooked meat samples, after the cooking loss assessment (AMSA, 1995); the same procedure was carried out on raw meat samples. The second subsample (weight 200 g) was used for proximate analysis; moisture and ash content were determined in ground freeze-dried samples. Crude protein was determined using a Kjeldahl Nitrogen/Protein Analyzer (Gerhardt Vapodest 50, Gerhardt. GMBH; Germany). The ash content was determined by incinerating samples in a muffle furnace at 500 °C. The LT fat content was determined with the Soxhlet method (2055 Soxtec Avanti, Foss Tecator AB, Höganäs, Sweden), according to the AOAC (1992). Intramuscular fat was extracted from LT muscle according to Folch et al. (1957). Fatty acid methyl esters were prepared according to the method proposed by Chistopherson et al. (1969). The upper organic phase was collected and filtered with anhydrous sodium sulphate and 1 µL was injected into a Fisons HRGC 8560 series MEGA2 gas chromatograph (Thermo Electron Corporation, Milan, Italy) equipped with a flame ionization detector and an AS 2000 automatic injection system (Thermo Electron Corporation, Milan, Italy). The column was a Varian (Varian Inc. Ass 1985, USA) CP-SIL 88 WCOT Fused Silica capillary column (100 m x 0.25 mm i.d. x 0.20 µm film thickness). FAME identification was performed by comparing their retention times with those of known standard FAME mixtures (SUPELCO, Sigma Aldrich Inc., St. Louis, MO). Each FAME was expressed as a percentage of the total areas of all FAME peaks. Peroxide value (PV) was determined in the lipid extract (50 mg) from the third subsample, as suggested by Shantha and Decker (1994). This method is based on the ability of peroxides to oxidize ferrous ions to ferric ones. Ammonium thiocyanate then reacts with ferric ions, resulting in a coloured complex that is measured at 500 nm with a spectrophotometer (double-beam UV-visible Jasco model V-550, Jasco International, Tokyo, Japan). PV was calculated from the

absorbance at 500 nm. For the quantitative determination of PV, an Fe(III) standard calibration curve was used with a concentration range of 0.1-5 $\mu\text{g/mL}$ ($y = 0.0282x - 0.0003$; $r^2 = 0.999$). PV was expressed as meq of O_2 per kg of fat. Three replicates were run per sample. TBA-reactive substances (TBARs) were determined in a 2 g sample (ground meat), according to the Witte (1970) method. This method is based on the reaction between thiobarbituric acid and the aldehydes that derive from secondary oxidation of the lipids present in the sample, resulting in a coloured complex that can be measured at 530 nm. TBARs were evaluated at such wavelength with a double-beam UV-visible spectrophotometer (Jasco model V-550, Jasco International, Tokyo, Japan), and they were calculated from the absorbance. For the quantitative determination of TBARs, a 1,1,3,3-tetramethoxypropane standard calibration curve was used with a concentration range of 0.045-0.113 $\mu\text{g/mL}$ ($y = 0.0087x - 0.0051$; $r^2 = 0.999$). The TBARs values were expressed as mg of malonylaldehyde (MDA) per kg of sample. Three replicates were run per sample. All data were analysed by one-way factorial ANOVA with repeated measures, using Tukey's HSD as *post-hoc* test. The dietary treatment was used as the source of variation to determine whether there were any significant differences ($P \leq 0.05$). Pearson's correlation coefficient ($\alpha = 0.05$) was used to separate the means of statistically different parameters and interactions and examine the possible relationships between the degree of unsaturation of the meat and the main oxidative indices. The Statistica version 10 software (StatSoft Inc., Padova, Italy) was used.

Results and discussion

Table 2 shows the chemical composition of the LT muscle: no significant increase in the lipid content of meat from heifers that had received flaxseed, or CLA, was noted. This result is in agreement with Juarez et al. (2012), who administered 10% of ground flaxseed to feedlot steers for 129 days. WBSF data here obtained, can be classified as "very tender", according to the Destefanis et al. (2008) WBSF scale (category n. 5; <3.36 kg). As observed by Juarez et al. (2012) and Nassu et al. (2011b), meat tenderness was not affected by lipid supplementation. Regarding the colour

1
2
3 131 measurements performed in the aged meat, L value (lightness) of group C was significantly less
4
5 132 bright than those of B, F and G groups. This finding might be related to those presented by Priolo et
6
7 133 al. (2001) who have noted by the linear regression between objective L value and time on pasture
8
9 134 that was significant ($P < 0.001$; $r^2 = 0.74$). The cause of this effect is extremely difficult to evaluate
10
11 135 because several factors play an important role, such as: carcass fatness, meat ultimate pH, animal
12
13 136 age, carcass weight and intramuscular fat content. Table 3 shows LT muscle fatty acid profile, in
14
15 137 particular for α -linolenic acid content, we observed a dose effect in groups supplemented with EF
16
17 138 compared with the A diet (C, D and E, average 0.31%). Moreover the α -linolenic acid content in a
18
19 139 short term EF supplementation (90 days, C, average 0.71%) is to prefer than a long term
20
21 140 supplementation (180 days, B, F and G, average 0.54%). Furthermore, Pearson test ($\alpha=0.05$),
22
23 141 between EF increasing doses (0, 250 and 500 g/head/day) and α -linolenic acid level in lipid meat,
24
25 142 was statistically significant ($r = 0.74825$). Priolo et al. (2001) reported the variation of α -linolenic
26
27 143 acid in intramuscular fat of cattle: grazing for six months on pasture has an effect of increasing
28
29 144 intramuscular α -linolenic acid by 50%. However, considering the lipid meat content shown in Table
30
31 145 3, these differences were relatively low from a quantitative point of view. Considering these results,
32
33 146 administering the same amount of flaxseed in a shorter time may be the best strategy (Nassu et al.,
34
35 147 2011a) to reduce partially the n-6/n-3 ratio (calculated on all FA detected) as noticed in previous
36
37 148 studies (Mordenti et al., 2005; Mordenti et al., 2013). Regarding the two main CLAs contents, our
38
39 149 results indicate that administering rp CLA, with or without EF, do not have had effectiveness on
40
41 150 both CLAs contents as compared to A group; obviously the CLAs contents were not influenced by
42
43 151 the treatment length. This finding is in agreement with what has been reported by other authors
44
45 152 (Schiavon et al., 2011) and might have been due to an CLA utilization directly by animals, reducing
46
47 153 body mass (trans-10, cis-12 CLA) and improving feed efficiency (cis-9, trans-11 CLA). Among
48
49 154 PUFA, arachidonic acid content was significantly higher ($P \leq 0.05$) in group A than in groups B and
50
51 155 D. The content of eicosatrienoic acid (C20:3 n-3 and n-6) was significantly higher ($P \leq 0.05$) in
52
53 156 group A than in groups B, D and G, and the A group amount of n-3 docosapentaenoic acid (DPA)

was significantly higher compared to the B. The n-6/n-3 ratio was significantly higher in group A than in groups B, C, F and G, by 27%, 37%, 25% and 30%, respectively. Regarding the oxidative parameters (Table 3), PV varied from 0.57 to 4.92 meq O₂/kg lipids, being significantly lower in the A diet with respect to the B diet. TBARs values ranged from 0.19 to 0.74 mg MDA/kg meat, remaining below the acceptance threshold of 2 mg MDA/kg beef meat (Campo et al., 2006). No significant effects of dietary supplementation on TBARs levels of fresh meat were observed. No correlations (Pearson test, $\alpha=0.05$) were found between the results obtained for lipid composition and the oxidative parameters resulting from the seven types of feeding conditions. In general, a low oxidation level was found in all samples, similar to that reported by Boselli et al. (2009), but both PV and TBARs were greater than those reported by other authors (Insani et al., 2008). This could be due to Italian carcass processing practices, which provide for a storage period, aimed at improving meat tenderness and promoting the formation of aroma compounds, which develop during cooking (Rodriguez-Estrada et al., 1997; Boselli et al., 2009). The long-term supplementation of EF seems to favour primary oxidation, even in the presence of vitamin E. Although the dietary supplementation did not have any significant impact on TBARs levels of fresh meat, the protective action of vitamin addition might become evident during meat storage under commercial retail conditions (Cardenia et al., 2011). On the other hand, it might be pointed out that TBARs describe an overall oxidative behaviour, which is useful if compared with PV; however, other specific antioxidant parameters (such as antioxidant enzyme activities and antioxidant capacity) could provide additional information (Mahecha et al., 2011).

Conclusions

The results obtained in the present study confirm that the diets that included EF supplementation proved to be effective at improving the n-3 FA and lowering the n-6/n-3 ratio in lipid fraction of beef meat. For this aspect, a shorter period of supplementation (90 days) was a more successful strategy than a longer one (180 days). However, considering the lipid meat content, these quantities

were relatively low for humans, from a quantitative point of view. Although oxidation parameters did not reflect the antioxidant effect of the latter, this effect might become evident during meat storage (Cardenia et al., 2011).

Disclosure statement

The authors report no conflicts of interest.

References

AOAC (Association of Official Analytical Chemists). 1992. (991.36) Fat (Crude) in Meat and Meat Products. Association of Official Analytical Chemists. Arlington, VA, USA.

AMSA. 1995. Research guidelines for cookery, sensory evaluation and instrumental tenderness measurements of fresh meat. In: American Meat Science Association and National Live Stock and Meat Board. Chicago, IL, USA.

Boselli E, Rodriguez-Estrada MT, Fedrizzi G, Caboni MF. 2009. Cholesterol photosensitised oxidation of beef meat under standard and modified atmosphere at retail conditions. Meat Sci. 81, 224-229.

Campo MN, Nute GR, Hughes SI, Enser M, Wood JD, Richardson RI. 2006. Flavour perception of oxidation in beef. Meat Sci. 72, 303–311.

Cardenia V, Rodriguez-Estrada MT, Cumella F, Sardi L, Della Casa G, Lercker G. 2011. Oxidative stability of pork meat lipids as related to high-oleic sunflower oil and vitamin E diet supplementation and storage conditions. Meat Sci. 88, 271-279.

Chistopherson S, Glass RL. 1969. Preparation of milk fat methyl esters by alcoholysis in an essential nonalcolic solution. J. Dairy Sci. 52, 1289-1290.

Destefanis G, Brugiapaglia A, Barge MT, Dal Molin E. 2008. Relationship between beef consumer tenderness perception and Warner–Bratzler shear force. Meat Sci. 78, 153–156.

- EFSA 2005. Opinion of the scientific panel on dietetic products, nutrition and allergies on a request from the commission related to nutrition claims concerning omega-3 fatty acids, monounsaturated fat, polyunsaturated fat and unsaturated fat. The EFSA Journal 253, 1-29.
- Folch J, Lees M, Sloane-Stanley GH. 1957. A simple method for the isolation and purification of total lipids from animal tissues. Journal of Biological Chemistry 226, 497–509.
- Honikel KO. 1998. Reference methods for the assessment of physical characteristics of meat. Meat Sci. 49, 447–457.
- Juarez M, Dugan MER, Aldai N, Basarab JA, Baron VS, McAllister TA, Aalhus JL. 2012. Beef quality attributes as affected by increasing the intramuscular levels of vitamin E and omega-3 fatty acids Meat Sci. 90, 764–769.
- Insani EM, Eyherabide A, Grigioni G, Sancho AM, Pensel NA, Descalzo AM. 2008. Oxidative stability and its relationship with natural antioxidants during refrigerated retail display of beef produced in Argentina. Meat Sci. 79, 444-452.
- Mahecha L, Nuernberg K, Nuernberg G, Martin J, Hubbermann EM, Knoeller S, Claeys E, De Smet S, Dannenberger D. 2011. Antioxidant enzyme activities an antioxidant capacity in *longissimus* muscle from bulls fed diets rich in polyunsaturated fatty acids. Food Chemistry 127, 379-386.
- McAfee AJ, McSorley EM, Cuskelly GJ, Moss BW, Wallace JMW, Bonham MP, Fearon AM. 2010. Red meat consumption: An overview of the risks and benefits. Meat Sci. 84,1–13.
- McNeill SH. 2014. Inclusion of red meat in healthful dietary patterns. Meat Sci. 98(3), 452-460
- Moloney AP, Scollan ND, Miles L. 2008. Enrichment of n-3 fatty acids and conjugated linoleic acid in beef: ProSafeBeef. Nutrition Bulletin 33, 374–381.
- Mordenti AL, Gatta PP, Lambertini L, Tassinari M. 2005. Effect of dietary supplementation with LCPUFA of vegetable origin in lipid content of muscular and subcutaneous tissues in heifers. Progress in Nutrition 7, 201-210.

- 233 Mordenti AL, Merendi F, Vecchia P, Panciroli A, Tassinari M, Brogna N, 2013. Effect of different
234 lipid supplements on steer performance, meat quality and fatty acid composition of Longissimus
235 dorsi muscle. ASPA 20th Congress Bologna, June 11-13. Ital. J. Anim. Sci. 12, C-036.
- 236 Nassu RT, Dugan MER, He ML, McAllister TA, Aalhus JL, Aldai JL, Kramer JKG. 2011a. The
237 effects of feeding flaxseed to beef cows given forage based diets on fatty acids of longissimus
238 thoracis muscle and backfat. Meat Sci. 89, 469-477.
- 239 Nassu RT, Dugan MER, Juarez M, Basarab JA, Baron VS, Aalhus JL, 2011b. Effect of α -
240 tocopherol tissue levels on beef quality. Animal 5, 2010-2018.
- 241 Priolo A, Micol D, Agabriel J. 2001. Effects of grass feeding systems on ruminant meat colour and
242 flavour. A review. Animal Research 50, 185-200.
- 243 Pouzo LB, Descalzo AM, Zaritzky NE, Rossetti L, Pavan E. 2016. Antioxidant status, lipid and
244 color stability of aged beef from grazing steers supplemented with corn grain and increasing
245 levels of flaxseed. Meat Sci. 111, 1-8.
- 246 Rodriguez-Estrada MT, Penazzi G, Caboni MF, Bertacco G, Lercker G. 1997. Effect of different
247 cooking methods on some lipid and protein components of hamburgers. Meat Sci. 45, 365-375.
- 248 Salter AM. 2013. Dietary fatty acids and cardiovascular disease. Animal 7, 163-171.
- 249 Schiavon S, Tagliapietra F, Dal Maso M, Bailoni L, Bittante G. 2011. Effects of low-protein diets
250 and rumen-protected conjugated linoleic acid on production and carcass traits of growing
251 double-muscled Piemontese bulls. Journal of Animal Science 88, 3372-3383.
- 252 Scollan ND, Dannenberger D, Nuenberg K, Richardson I, MacKintosh S, Hocquette JF, Moloney
253 AP. 2014. Enhancing the nutritional and health value of beef lipids and their relationship with
254 quality. Meat Sci. 97, 384-394.
- 255 Shantha NC, Decker EA. 1994. Rapid, sensitive, iron-based spectrophotometric method for
256 determination of peroxide values of food lipids. Journal of AOAC International 77, 421-424.
- 257 U.S. Department of Health and Human Services. 2010. Dietary guidelines for Americans. Retrieved
258 September 17, 2014, from. <http://health.gov/dietaryguidelines/2010.asp>

- 1
2
3 259 Van Soest PJ, Robertson JB, Lewis BA. 1991. Methods for dietary fiber neutral detergent fiber, and
4
5 260 non-starch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74, 583-597.
6
7 261 Witte VC, Krause GF, Bailet ME. 1970. A new extraction method for determining 2-thiobarbituric
8
9 262 acid values of pork and beef during storage. *J. Food Sci.* 35, 582–585.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

1 **Tables**

2 Table 1: Description of the seven dietary treatments administered to heifers (g as feed)

Groups	Supplementation	Lenght	Description
A (control)		180	Basal diet supplemented with Megalac® and soybean meal (protein 14% and energy 1762 kcal/kg)
B	EF/VE	180	Basal diet plus 250 g EF head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
C	Control + EF/VE	90 + 90	Basal diet (first 90 days) followed by basal diet supplemented with 500 g EF head ⁻¹ day ⁻¹ and 2 g VE head ⁻¹ day ⁻¹
D	CLA/VE	180	Basal diet supplemented with 25 g CLA head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
E	Control + CLA/VE	90 + 90	Basal diet (first 90 days) followed by basal diet supplemented with 50 g CLA head ⁻¹ day ⁻¹ and 2 g VE head ⁻¹ day ⁻¹
F	EF/CLA/VE	180	Basal diet supplemented with 250 g EF head ⁻¹ day ⁻¹ , 25 g CLA head ⁻¹ day ⁻¹ and 1 g VE head ⁻¹ day ⁻¹
G	EF/CLA	180	Basal diet supplemented with 250 g EF head ⁻¹ day ⁻¹ and 25 g CLA head ⁻¹ day ⁻¹

3 EF, extruded flaxseed; VE, DL-α tocopheril acetate; CLA, conjugated linoleic acid isomers.

Table 2: Composition (% of DM) and physical properties of the *Longissimus* muscle in heifers

Item	Experimental groups							SEM	P
	A	B	C	D	E	F	G		
Dry Matter	26.9	28.1	26.8	27.2	26.6	27.8	27.2	0.15	ns
Protein	79.0	77.6	78.2	79.3	80.2	78.3	80.3	0.57	ns
Ether extract	13.6	15.3	14.6	12.7	11.9	14.1	12.7	0.56	ns
Cooking loss (%)	30.1	28.9	30.3	30.2	30.2	28.3	29.1	0.37	ns
WBSF fresh (kg)	1.84	1.75	2.10	2.08	2.08	1.89	1.76	0.06	ns
WBSF cooked (kg)	3.38	3.52	3.55	3.23	3.77	4.29	3.56	0.10	ns
Color at day 14									
L value	42.7 ^{ab}	44.6 ^a	40.9 ^b	42.9 ^{ab}	43.3 ^{ab}	44.5 ^a	45.4 ^a	0.35	≤ 0.05
a* value	20.9	21.8	21.3	20.5	20.7	20.2	20.0	0.23	ns
b* value	11.0	11.8	10.5	10.4	10.8	10.6	11.2	0.19	ns

Values in the same row with different superscripts are different (a and b, $P \leq 0.05$; A and B $P \leq 0.01$).

SEM=standard error of the mean; WBSF = Warner-Bratzler Shear Force

Table 3: Fatty acid content, fatty acid classes and oxidative parameters of *Longissimus* muscle in heifers

Fatty acids	Experimental groups							SEM	P
(% of total FA)	A	B	C	D	E	F	G		
C14:0	2.95	3.05	3.18	3.33	3.24	2.95	2.80	0.08	ns
C16:0	28.34	28.61	28.22	29.32	28.51	28.54	27.76	0.25	ns
C16:1 n-7	2.67	2.55	2.45	2.57	2.47	2.74	2.45	0.05	ns
C18:0	17.43	19.22	19.67	19.54	19.51	17.44	19.21	0.26	ns
C18:1 n-9	35.92	36.17	33.59	34.83	34.45	35.97	37.48	0.36	ns
C18:1 n-7	1.94 ^{ab}	1.59 ^{ab}	1.99 ^a	1.65 ^{ab}	1.84 ^{ab}	1.68 ^{ab}	1.43 ^b	0.14	≤ 0.05
C18:2 n-6	6.20	4.91	6.04	5.00	5.59	6.29	4.86	0.21	ns
C18:3 n-3	0.29 ^{Bb}	0.54 ^{Ac}	0.71 ^{Aa}	0.33 ^{Bb}	0.30 ^{Bb}	0.56 ^{Ac}	0.52 ^{Ac}	0.02	≤ 0.05
CLA cis9,trans11	0.34	0.41	0.40	0.33	0.35	0.34	0.36	0.03	ns
CLA trans10,cis12	0.16	0.15	0.14	0.13	0.12	0.15	0.16	0.01	ns
C20:4 n-6	1.10 ^a	0.48 ^b	0.93 ^{ab}	0.57 ^b	0.88 ^{ab}	0.83 ^{ab}	0.58 ^b	0.05	≤ 0.05
C20:5 n-3	0.10 ^{ab}	0.06 ^b	0.12 ^a	0.07 ^b	0.08 ^{ab}	0.11 ^a	0.06 ^b	0.01	≤ 0.05
C22:5 n-3	0.14 ^A	0.06 ^B	0.11 ^{AB}	0.09 ^{AB}	0.11 ^{AB}	0.11 ^{AB}	0.09 ^{AB}	0.01	≤ 0.01
C22:6 n-3	0.29	0.16	0.29	0.18	0.23	0.29	0.19	0.02	ns
SFA	50.01 ^B	52.26 ^{AB}	52.51 ^A	53.61 ^A	52.83 ^A	50.21 ^B	51.14 ^B	0.83	≤ 0.01
MUFA	41.08	40.80	38.53	39.53	39.29	40.89	41.88	0.89	ns
PUFA	8.91	6.94	8.96	6.86	7.88	8.89	6.98	0.74	ns
PUFA/SFA	1.00 ^A	0.91 ^{AB}	0.91 ^{AB}	0.87 ^B	0.90 ^{AB}	1.00 ^{AB}	0.96 ^{AB}	0.03	≤ 0.01
n-6/n-3	8.99 ^a	6.54 ^b	5.70 ^b	8.63 ^a	8.98 ^a	6.76 ^b	6.26 ^b	0.46	≤ 0.05
TBARs	0.70	0.26	0.19	0.44	0.21	0.32	0.73	0.23	ns
PV	0.54 ^b	4.11 ^a	0.85 ^b	2.80 ^{ab}	1.07 ^{ab}	4.28 ^a	2.59 ^{ab}	1.45	≤ 0.05

Values in the same row with different superscripts are different (a and b, P≤0.05; A and B P≤0.01).

1 SEM= standard error of the mean; SFA = saturated fatty acids; MUFA = monounsaturated fatty
2 acids; PUFA = polyunsaturated fatty acids; TBARs = TBA-reactive substances (mg MDA/kg
3 meat); PV = peroxide value (meq O₂/kg lipids)
4

For Peer Review Only